

Southeast Ag Solutions

N-Terragate LH



Microbial Soil and Foliar Drench Amendment

MICROBIAL INNOCULANTS 16%

Bacillus subtilis 1538.....	5.0 x 10 ⁹ cfu/g
Bacillus amyloliquefaciens 1147.....	5.0 x 10 ⁹ cfu/g
Bacillus subtilis 1952.....	2.5 x 10 ⁹ cfu/g
Bacillus plantarum 1595.....	2.5 x 10 ⁹ cfu/g

OTHER INGREDIENTS 84%

Microbial Nutrients-Carbohydrates

TOTAL 100%

* Contains a combined minimum cfu (colony forming unit) of 1.5x10¹⁰ per gram

Packaging:

- 3 LB Container
- 4x3 LB Containers
- 40 LB Container

Manufactured for:

Southeast Ag
Solutions
P.O. Box 7964
Sebring, FL 33872
863-382-2737

Manufactured by:

Asymmetry Technologies
104 Three West Parkway
#300
Villa Rica, GA 30180
855-553-2716

SIGNAL WORD: This product contains no substance which, at their given concentration, are considered to be hazardous to health.

WARNING HAZARD STATEMENTS: None

PRECAUTIONARY STATEMENTS – PREVENTION: As a precaution, wear chemical resistance gloves (PVC, neoprene or butyl). Safety glasses or chemical goggles are recommended.

PRECAUTIONARY STATEMENTS – RESPONSE:

If swallowed: Rinse mouth and throat. Drink water to dilute.

If on skin (or hair): Rinse skin with and non-abrasive soap. Remove contaminated clothing and wash before reuse. If skin irritation occurs, get medical advice/attention.

If inhaled: Remove person to fresh air and keep comfortable for breathing.

If in eyes: Rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice/attention.

STORAGE AND HANDLING: Store in a cool, dry place. Avoid contact with strong acids or bases.

FIRE: This product is non-combustible. No extinguishing media or special fire-fighting procedures are required. There are no unusual fire or explosion hazards.

DISPOSAL: Dispose of in licensed landfills or incinerate in accordance with local, state and federal regulations.

DIRECTIONS FOR USE: (8 ounces = 1 cup)

Recommended treatment is through Chemigation (drip, trickle or micro-jet irrigation), Foliar Spray or Drench. A stock tank may be used to dose product. Partially fill the mix tank with clean water. Add the specified amount of N-Terragate LH to the tank. Finish filling the tank with clean water to the necessary volume required. The solution should be agitated during mixing and application to assure a uniform suspension. N-Terragate LH can be mixed and used with other agricultural chemicals, for which such mixing is permitted by the product label. If such a mixture is planned, a compatibility test should be first conducted. Avoid mixing with quaternary ammonia compounds, Peroxyacetic Acid (PAA), or similar type broad spectrum biocides. Do not allow mixture to stand overnight or for prolonged periods. Mix well before each use. Dose 4-12 ounces per 100 gallons of water. For specific dosage rates and frequency of treatment, see Dosage Table on Product Data Sheet or contact your Account Representative.

Refer to product Safety Data Sheet for further safety and health information. For product information, call 863-382-2737.