



N-Terragate T2 +B PM

Microbial Soil and Foliar Drench Amendment

MICROBIAL INNOCULANTS.....	16%
Trichoderma <i>harzianum</i>	2.5 x 10 ⁸ cfu/g
Trichoderma <i>viride</i>	2.5 x 10 ⁸ cfu/g
Bacillus blend.....	1.0 x 10 ⁸ cfu/g
(Bacillus blend= Bacillus <i>subtilis</i> , Bacillus <i>licheniformis</i> , Bacillus <i>megaterium</i>)	
OTHER INGREDIENTS.....	84%
Microbial Nutrients-Carbohydrates	
TOTAL.....	100%
* Contains a combined minimum cfu (colony forming unit) of 6.0x10 ⁸ per gram	

IT IS THE USERS RESPONSIBILITY TO DETERMINE CONDITIONS FOR THE USE OF THIS PRODUCT AND TO ASSUME LIABILITY FOR LOSS, INJURY, DAMAGE OR EXPENSE RESULTUING IN IMPROPER USE.

Refer to product Safety Data Sheet for further safety and health information

PACKAGING: 25LB containers.

SIGNAL WORD: This product contains no substance which, at their given concentration, are considered to be hazardous to health.

WARNING HAZARD STATEMENTS: None

PRECAUTIONARY STATEMENTS - PREVENTION: As a precaution, wear chemical resistance gloves (PVC, neoprene or butyl). Safety glasses or chemical goggles are recommended.

PRECAUTIONARY STATEMENTS - RESPONSE:

If swallowed: Rinse mouth and throat. Drink water to dilute.

If on skin (or hair): Rinse skin with and non-abrasiive soap. Remove contaminated clothing and wash before reuse. If skin irritation occurs, get medical advice/attention.

If inhaled: Remove person to fresh air and keep comfortable for breathing.

If in eyes: Rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice/attention.

STORAGE AND HANDLING: Do not ingest internally. Avoid skin contact. Wear rubber gloves and eye protection when using this product. Wash hands thoroughly after handling.

FIRE: This product is non-combustible. No extinguishing media or special fire-fighting procedures are required. There are no unusual fire or explosion hazards.

DISPOSAL: Dispose of in licensed landfills or incinerate in accordance with local, state and federal regulations.

DIRECTIONS FOR USE: (8 ounces=1 cup)

Recommended treatment is through Chemigation (drip, trickle or microjet irrigation), Foliar Spray or Drench. A stock tank may be used to dose product. Partially fill the mix tank with clean water. Add the specified amount of **N-Terragate T2+B PM** to the tank. Finish filling the tank with clean water to the necessary volume. The solution should be agitated during mixing and application to assure uniform suspension. **N-Terragate T2+B PM** can be mixed and used with other agricultural chemicals, for which such mixing is permitted by the product label. If such a mixture is planned, a compatibility test should be first conducted. Avoid mixing with quaternary compounds, Peroxyacetic Acid (PAA), or similar type broad spectrum biocides. Do not allow mixture to stand overnight or for prolonged periods. Mix well before each use. Dose 4-12 ounces per 100 gallons of water. For specific dosage rates and frequency of treatment, contact your Account Representative.

Southeast Ag Solutions, Inc. P.O. Box 7964. Sebring, FL 33872. 863-382-2737